

Detailed Steps of the Fourier-Motzkin Elimination

Xiaoli Xu, Yong Zeng, Yong Liang Guan and Tracey Ho

This file provide the detailed steps for obtaining the bounds on R_1 , R_2 via the obtained results on $(R_{1c}, R_{1p}, R_{2c}, R_{2p})$. The rate pair $(R_{1c}, R_{1p}, R_{2c}, R_{2p})$ is achievable if it satisfies following inequalities.

$$R_{1c} \leq \text{rank}(U_1^T H_{11} V_{21}) \quad (1)$$

$$R_{1p} \leq m_1 - r_{21} \quad (2)$$

$$R_{2c} \leq r_{12} \quad (3)$$

$$R_{1c} + R_{1p} \leq r_{11} \quad (4)$$

$$R_{1c} + R_{2c} \leq r_{12} + \text{rank}(U_{10}^T H_{11} V_{21}) \quad (5)$$

$$R_{1p} + R_{2c} \leq r_{12} + \text{rank}(U_{10}^T H_{11} V_{20}) \quad (6)$$

$$R_{1c} + R_{1p} + R_{2c} \leq n_1 \quad (7)$$

$$R_{2c} \leq \text{rank}(U_2^T H_{22} V_{11}) \quad (8)$$

$$R_{2p} \leq m_2 - r_{12} \quad (9)$$

$$R_{1c} \leq r_{21} \quad (10)$$

$$R_{2c} + R_{2p} \leq r_{22} \quad (11)$$

$$R_{2c} + R_{1c} \leq r_{21} + \text{rank}(U_{20}^T H_{22} V_{11}) \quad (12)$$

$$R_{2p} + R_{1c} \leq r_{21} + \text{rank}(U_{20}^T H_{22} V_{10}) \quad (13)$$

$$R_{2c} + R_{2p} + R_{1c} \leq n_2 \quad (14)$$

where (3) is implied by (8) as $\text{rank}(U_2^T H_{22} V_{11}) \leq \text{rank}(V_{11}) = r_{12}$. Similarly, (10) is implied by (1). Removing the redundant inequalities and substituting with $R_{1p} = R_1 - R_{1c}$, $R_{2p} = R_2 - R_{2c}$,

we obtain

$$R_{1c} \leq \text{rank}(U_1^T H_{11} V_{21}) \quad (15)$$

$$R_1 - R_{1c} \leq m_1 - r_{21} \quad (16)$$

$$R_1 \leq r_{11} \quad (17)$$

$$R_{1c} + R_{2c} \leq r_{12} + \text{rank}(U_{10}^T H_{11} V_{21}) \quad (18)$$

$$R_1 - R_{1c} + R_{2c} \leq r_{12} + \text{rank}(U_{10}^T H_{11} V_{20}) \quad (19)$$

$$R_1 + R_{2c} \leq n_1 \quad (20)$$

$$R_{2c} \leq \text{rank}(U_2^T H_{22} V_{11}) \quad (21)$$

$$R_2 - R_{2c} \leq m_2 - r_{12} \quad (22)$$

$$R_2 \leq r_{22} \quad (23)$$

$$R_{2c} + R_{1c} \leq r_{21} + \text{rank}(U_{20}^T H_{22} V_{11}) \quad (24)$$

$$R_2 - R_{2c} + R_{1c} \leq r_{21} + \text{rank}(U_{20}^T H_{22} V_{10}) \quad (25)$$

$$R_2 + R_{1c} \leq n_2 \quad (26)$$

To eliminate R_{1c} , we get five upper bounds and two lower bounds on R_{1c}

$$R_{1c} \leq \text{rank}(U_1^T H_{11} V_{21})$$

$$R_{1c} \leq r_{12} + \text{rank}(U_{10}^T H_{11} V_{21}) - R_{2c}$$

$$R_{1c} \leq r_{21} + \text{rank}(U_{20}^T H_{22} V_{11}) - R_{2c}$$

$$R_{1c} \leq r_{21} + \text{rank}(U_{20}^T H_{22} V_{10}) - R_2 + R_{2c}$$

$$R_{1c} \leq n_2 - R_2$$

$$R_{1c} \geq R_1 - m_1 + r_{21}$$

$$R_{1c} \geq R_1 + R_{2c} - r_{12} - \text{rank}(U_{10}^T H_{11} V_{20})$$

Comparing the upper and lower bounds of R_{1c} and copying five inequalities in (15)-(26) that do

not involve R_{1c} , we obtain a new system of inequalities in (R_{2c}, R_1, R_2) give by

$$R_1 \leq r_{11} \quad (27)$$

$$R_1 + R_{2c} \leq n_1 \quad (28)$$

$$R_{2c} \leq \text{rank}(U_2^T H_{22} V_{11}) \quad (29)$$

$$R_2 - R_{2c} \leq m_2 - r_{12} \quad (30)$$

$$R_2 \leq r_{22} \quad (31)$$

$$R_1 \leq \text{rank}(U_1^T H_{11} V_{21}) + m_1 - r_{21} \quad (32)$$

$$R_1 + R_{2c} \leq r_{12} + \text{rank}(U_{10}^T H_{11} V_{21}) + m_1 - r_{21} \quad (33)$$

$$R_1 + R_{2c} \leq \text{rank}(U_{20}^T H_{22} V_{11}) + m_1 \quad (34)$$

$$R_1 + R_2 - R_{2c} \leq \text{rank}(U_{20}^T H_{22} V_{10}) + m_1 \quad (35)$$

$$R_1 + R_2 \leq n_2 + m_1 - r_{21} \quad (36)$$

$$R_1 + R_{2c} \leq \text{rank}(U_1^T H_{11} V_{21}) + \text{rank}(U_{10}^T H_{11} V_{20}) + r_{12} \quad (37)$$

$$R_1 + 2R_{2c} \leq 2r_{12} + \text{rank}(U_{10}^T H_{11} V_{21}) + \text{rank}(U_{10}^T H_{11} V_{20}) \quad (38)$$

$$R_1 + 2R_{2c} \leq r_{21} + \text{rank}(U_{20}^T H_{22} V_{11}) + \text{rank}(U_{10}^T H_{11} V_{20}) + r_{12} \quad (39)$$

$$R_1 + R_2 \leq \text{rank}(U_{20}^T H_{22} V_{10}) + \text{rank}(U_{10}^T H_{11} V_{20}) + r_{12} + r_{21} \quad (40)$$

$$R_1 + R_2 + R_{2c} \leq n_2 + r_{12} + \text{rank}(U_{10}^T H_{11} V_{20}) \quad (41)$$

where (32) is implied by (27) as $\text{rank}(U_1^T H_{11} V_{21}) = \text{rank}(H_{11} V_{21}) \geq r_{11} - m_1 + r_{21}$; (33) is implied by (28) as $\text{rank}(U_{10}^T H_{11} V_{21}) \geq n_1 - m_1 + r_{21} - r_{12}$; (37) is implied by (28) as $\text{rank}(U_1^T H_{11} V_{21}) + \text{rank}(U_{10}^T H_{11} V_{20}) + r_{12} = \text{rank}(H_{11} V_{21}) + \text{rank}(U_{10}^T H_{11} V_{20}) + r_{12} \geq r_{11} + r_{21} - m_1 + r_{12} + n_1 + m_1 - r_{11} - r_{12} - r_{21} = n_1$.

Now, we are ready to eliminate R_{2c} from the remaining non-redundant inequalities. In total,

six upper bounds and two lower bounds are obtained for R_{2c} .

$$R_{2c} \leq n_1 - R_1$$

$$R_{2c} \leq \text{rank}(U_2^T H_{22} V_{11})$$

$$R_{2c} \leq \text{rank}(U_{20}^T H_{22} V_{11}) + m_1 - R_1$$

$$2R_{2c} \leq r_{21} + \text{rank}(U_{20}^T H_{22} V_{11}) + \text{rank}(U_{10}^T H_{11} V_{20}) + r_{12} - R_1$$

$$R_{2c} \leq n_2 + r_{12} + \text{rank}(U_{10}^T H_{11} V_{20}) - R_1 - R_2$$

$$2R_{2c} \leq 2r_{12} + \text{rank}(U_{10}^T H_{11} V_{21}) + \text{rank}(U_{10}^T H_{11} V_{20}) - R_1$$

$$R_{2c} \geq R_2 - m_2 + r_{12}$$

$$R_{2c} \geq R_1 + R_2 - \text{rank}(U_{20}^T H_{22} V_{10}) - m_1$$

Comparing the lower and upper bounds of R_{2c} , we get 12 inequalities. Together with the inequalities in (27)-(38) that do not involve R_{2c} , we obtain a new system of inequalities in

(R_1, R_2) given by

$$R_1 \leq r_{11} \quad (42)$$

$$R_2 \leq r_{22} \quad (43)$$

$$R_1 + R_2 \leq n_2 + m_1 - r_{21} \quad (44)$$

$$R_1 + R_2 \leq \text{rank}(U_{20}^T H_{22} V_{10}) + \text{rank}(U_{10}^T H_{11} V_{20}) + r_{12} + r_{21} \quad (45)$$

$$R_1 + R_2 \leq n_1 + m_2 - r_{12} \quad (46)$$

$$R_2 \leq \text{rank}(U_2^T H_{22} V_{11}) + m_2 - r_{12} \quad (47)$$

$$R_1 + R_2 \leq \text{rank}(U_{20}^T H_{22} V_{11}) + m_1 + m_2 - r_{12} \quad (48)$$

$$R_1 + 2R_2 \leq r_{21} + \text{rank}(U_{20}^T H_{22} V_{11}) + \text{rank}(U_{10}^T H_{11} V_{20}) + 2m_2 - r_{12} \quad (49)$$

$$R_1 + 2R_2 \leq n_2 + m_2 + \text{rank}(U_{10}^T H_{11} V_{20}) \quad (50)$$

$$R_1 + 2R_2 \leq \text{rank}(U_{10}^T H_{11} V_{21}) + \text{rank}(U_{10}^T H_{11} V_{20}) + 2m_2 \quad (51)$$

$$2R_1 + R_2 \leq n_1 + \text{rank}(U_{20}^T H_{22} V_{10}) + m_1 \quad (52)$$

$$R_1 + R_2 \leq \text{rank}(U_2^T H_{22} V_{11}) + \text{rank}(U_{20}^T H_{22} V_{10}) + m_1 \quad (53)$$

$$2R_1 + R_2 \leq \text{rank}(U_{20}^T H_{22} V_{11}) + 2m_1 + \text{rank}(U_{20}^T H_{22} V_{10}) \quad (54)$$

$$3R_1 + 2R_2 \leq \text{rank}(U_{20}^T H_{22} V_{11}) + r_{12} + r_{21} + \text{rank}(U_{10}^T H_{11} V_{20}) + 2\text{rank}(U_{20}^T H_{22} V_{10}) + 2m_1 \quad (55)$$

$$2R_1 + 2R_2 \leq n_2 + r_{12} + \text{rank}(U_{10}^T H_{11} V_{20}) + \text{rank}(U_{20}^T H_{22} V_{10}) + m_1 \quad (56)$$

$$3R_1 + 2R_2 \leq 2r_{12} + \text{rank}(U_{10}^T H_{11} V_{21}) + \text{rank}(U_{10}^T H_{11} V_{20}) + 2\text{rank}(U_{20}^T H_{22} V_{10}) + 2m_1 \quad (57)$$

Now, we are proceed to identifying the redundant inequalities.

(47) is implied by (43) as $\text{rank}(U_2^T H_{22} V_{11}) = \text{rank}(H_{22} V_{11}) \geq r_{22} + r_{12} - m_2$.

(49) is implied by (50) as $\text{rank}(U_{20}^T H_{22} V_{11}) \geq n_2 - m_2 + r_{12} - r_{21}$.

(51) is implied by (43) and (46) as

$$\begin{aligned} \text{rank}(U_{10}^T H_{11} V_{21}) + \text{rank}(U_{10}^T H_{11} V_{20}) + 2m_2 &\geq \text{rank}(U_{10}^T H_{11}) + 2m_2 \\ &\geq n_1 - r_{12} + 2m_2 \geq n_1 + m_2 - r_{12} + r_{22} \end{aligned}$$

(48) is implied by (44) as $\text{rank}(U_{20}^T H_{22} V_{11}) \geq n_2 - m_2 + r_{12} - r_{21}$.

(53) is implied by (44) as $\text{rank}(U_2^T H_{22} V_{11}) + \text{rank}(U_{20}^T H_{22} V_{10}) + m_1 = \text{rank}(H_{22} V_{11}) + \text{rank}(U_{20}^T H_{22} V_{10}) + m_1 \geq n_2 - r_{21} + m_1$.

(55) can be obtained by summing up (45) and (54).

(56) can be obtained by summing up (44) and (45).

(57) is implied by (45) and (52).

(54) is implied by (42) and (44) as

$$\begin{aligned} \text{rank}(U_{20}^T H_{22} V_{11}) + 2m_1 + \text{rank}(U_{20}^T H_{22} V_{10}) &\geq \text{rank}(U_{20}^T H_{22}) + 2m_1 \\ &= n_2 - r_{21} + 2m_1 \geq n_2 - r_{21} + m_1 + r_{11} \end{aligned}$$

After removing all the redundant inequalities, we obtain the non-trivial set of inequalities.

$$R_1 \leq r_{11} \quad (58)$$

$$R_2 \leq r_{22} \quad (59)$$

$$R_1 + R_2 \leq n_2 + m_1 - r_{21} \quad (60)$$

$$R_1 + R_2 \leq n_1 + m_2 - r_{12} \quad (61)$$

$$R_1 + R_2 \leq \text{rank}(U_{20}^T H_{22} V_{10}) + \text{rank}(U_{10}^T H_{11} V_{20}) + r_{21} + r_{12} \quad (62)$$

$$2R_1 + R_2 \leq \text{rank}(U_{20}^T H_{22} V_{10}) + m_1 + n_1 \quad (63)$$

$$R_1 + 2R_2 \leq \text{rank}(U_{10}^T H_{11} V_{20}) + m_2 + n_2 \quad (64)$$

APPENDIX A

LIST OF FACTS

List of facts that are useful for in Fourier-Motzkin Elimination

- 1) $\text{rank}(U_{10}^T H_{11}) = n_1 - r_{12}$
- 2) $\text{rank}(U_{20}^T H_{22}) = n_2 - r_{21}$
- 3) $\text{rank}(H_{11} V_{20}) = m_1 - r_{21}$
- 4) $\text{rank}(H_{22} V_{10}) = m_2 - r_{12}$
- 5) $\text{rank}(H_{11} V_{21}) \geq r_{11} + r_{21} - m_1$
- 6) $\text{rank}(H_{22} V_{11}) \geq r_{22} + r_{12} - m_2$
- 7) $\text{rank}(U_{10}^T H_{11} V_{21}) \geq n_1 - m_1 + r_{21} - r_{12}$
- 8) $\text{rank}(U_{20}^T H_{22} V_{11}) \geq n_2 - m_2 + r_{12} - r_{21}$

$$9) \text{ rank}(U_{10}^T H_{11} V_{20}) \geq n_1 + m_1 - r_{11} - r_{12} - r_{21}$$

$$10) \text{ rank}(U_{20}^T H_{22} V_{10}) \geq n_2 + m_2 - r_{22} - r_{12} - r_{21}$$

Proof: Fact 1 can be proved as

$$\begin{aligned} \text{rank}(U_{10}^T H_{11}) &= \text{rank}(H_{11}) - \dim(\mathcal{N}(U_{10}^T) \cap \mathcal{R}(H_{11})) \\ &= r_{11} - \dim(\mathcal{R}(H_{12}) \cap \mathcal{R}(H_{11})) \\ &= r_{11} - (r_{11} + r_{12} - \text{rank}([H_{11} \ H_{12}])) \\ &= \text{rank}([H_{11} \ H_{12}]) - r_{12} \\ &\stackrel{(a)}{=} n_1 - r_{12} \end{aligned}$$

where (a) follows as $[H_{11} \ H_{12}]$ is full row rank. By symmetry, Fact 2 can be proved.

To prove Fact 3

$$\begin{aligned} \text{rank}(H_{11} V_{20}) &= \text{rank}(V_{20}) - \dim(\mathcal{N}(H_{11}) \cap \mathcal{R}(V_{20})) \\ &= m_1 - r_{21} - \dim(\mathcal{N}(H_{11}) \cap \mathcal{N}(H_{21})) \\ &\stackrel{(b)}{=} m_1 - r_{21} \end{aligned}$$

where (b) follows as $\begin{bmatrix} H_{11} \\ H_{21} \end{bmatrix}$ is full column rank. By symmetry, Fact 4 can be proved.

Fact 5 can be proved as $\text{rank}(H_{11}) = \text{rank}\left(H_{11} \begin{bmatrix} V_{21} & V_{20} \end{bmatrix}\right) \leq \text{rank}(H_{11} V_{21}) + \text{rank}(H_{11} V_{20})$.

Since $\text{rank}(H_{11} V_{20}) = m_1 - r_{21}$, $\text{rank}(H_{11} V_{21}) \geq r_{11} + r_{21} - m_1$. Fact 6 follows similarly.

Fact 7 can be proved using Frobenius inequality,

$$\begin{aligned} \text{rank}(U_{10}^T H_{11} V_{21}) &\geq \text{rank}(U_{10}^T H_{11}) + \text{rank}(H_{11} V_{21}) - \text{rank}(H_{11}) \\ &\geq n_1 - r_{12} + r_{11} + r_{21} - m_1 - r_{11} \\ &= n_1 - m_1 + r_{21} - r_{12} \end{aligned}$$

Fact 8 follows similarly.

Fact 9 is also proved using Frobenius inequality,

$$\begin{aligned} \text{rank}(U_{10}^T H_{11} V_{20}) &\geq \text{rank}(U_{10}^T H_{11}) + \text{rank}(H_{11} V_{20}) - \text{rank}(H_{11}) \\ &= n_1 - r_{12} + m_1 - r_{21} - r_{22} \end{aligned}$$

Fact 10 follows similarly. ■